

Herbaceous Biomass Estimation from SPOT-5 Imagery in Semiarid Rangelands of Idaho

Fang Chen, GIS Training and Research Center, Idaho State University, 921 S. 8th Ave, Stop 8104, Pocatello Idaho 83209-8104, chenfang@isu.edu

Keith T. Weber, GIS Director, Idaho State University. GIS Training and Research Center, 921 S. 8th Ave., Stop 8104, Pocatello, ID 83209-8104. webekeit@isu.edu

Bhushan Gokhale, GIS Training and Research Center, Idaho State University, 921 S. 8th Ave, Stop 8104, Pocatello Idaho 83209-8104

ABSTRACT

Eight vegetation indices (VI) commonly used for above-ground biomass (AGB) estimation were derived from Satellite Pour l'Observation de la Terre 5 (SPOT 5) imagery and used to predict herbaceous AGB at a semiarid rangeland study site in southeast Idaho. The relationship between herbaceous AGB and vegetation water content was also evaluated and as a result, a suite of water sensitive vegetation indices (WSVI) were developed. Correlation coefficients between herbaceous AGB, VI's, and WSVI's were calculated, demonstrating that WSVI's were correlated ($r^2 \geq 0.51$) with vegetation water content and performed better than standard VI's in herbaceous AGB estimates within the semiarid rangelands of Idaho.

KEYWORDS: Rangelands, biomass, water content, vegetation indices, remote sensing, Idaho

INTRODUCTION

Rangelands cover approximately 40% of the earth’s terrestrial surface and are important areas for livestock production and wildlife habitat (Breman and de Wit 1983; Huntsinger and Hopkins 1996). To effectively manage rangelands it is important to assess ecosystem productivity and biomass production (Running et al. 2004). Biomass estimates represent the quantity of matter in a given area and are expressed either as the weight of organisms per unit area or as the volume of organisms per unit volume. Previous total above-ground biomass (AGB) research has demonstrated that vegetation indices (VI) are sensitive to the biophysical and biochemical variations in vegetation and as a result, are the most common parameters used to estimate AGB (Davidson and Csillag 2001; Kawamura et al. 2005; Numata et al. 2008). A remote sensing derived VI is a quantitative optical measure of canopy greenness (Tucker 1979; Weiser et al. 1986). Various VI's such as the normalized difference vegetation index (NDVI), normalized difference water index (NDWI) and soil adjusted vegetation index (SAVI) have been correlated with AGB, and applied to predict AGB within a variety of biomes (Davidson and Csillag 2001; Kogan et al. 2004; Mirik et al. 2005; Wessels et al. 2006; Numata et al. 2008; Cho and Skidmore 2009) (Table 1). Recently, ground-based and satellite-based spectral measurement methods have been developed to better quantify AGB. For instance, many ground-based methods use portable field spectroradiometers or digital cameras (e.g., ASD spectrometers, ASD Inc, Boulder, CO,USA; Dycam Agricultural Digital Camera (ADC), Dycam Inc, Chatworth, CA, USA) to collect canopy radiance and predict AGB through an empirical relationship between spectral values and biomass samples (Boelman et al. 2003; Flynn et al. 2008; Mašková et al. 2008). These methods are straightforward and accurate for small-scale studies (e.g., approximately 1-10 ha), however, they are also labor intensive and difficult to apply over broad spatial scales or long-term temporal scales.

Table 1. Vegetation indices correlation with total above ground biomass (AGB) reported in different studies

<i>Study area</i>	<i>Sensor</i>	<i>Index</i>	<i>R²</i>	<i>Sources</i>
Kentucky, USA	Greenseeker RT500	NDVI	0.68	Flynn et al., 2008
Southern Africa	Landsat7 -ETM+	Green/Blue	0.85	Samimi and Kraus, 2004
Inner Mongolia, China	MODIS	NDVI	0.75	Kawamura et al., 2005
Italy	HyMap	NDVI	0.32-0.58	Cho and Skidmore, 2009
		NDWI	0.49-0.55	
Czech Republic	ADC	NDVI	0.83 (managed) 0.52 (unmanaged)	Mašková et al., 2008
Namibia	AVHRR	NDVI	0.76	Sannier et al., 2002
Northern Alaska	UniSpec-DC	NDVI	0.84	Boelman et al., 2003
Italy		NDVI	0.32	Schino et al., 2003
Brazilian Amazon	Analytical Spectral Device	NDVI	0.03	Numata et al., 2008
		NDWI	0.13	

The increasing availability of satellite-based remote sensing data extends the assessment of AGB into a broader spatiotemporal scale. For example, remotely sensed data acquired from various sensors have been used to assess AGB including NOAA's AVHRR (Box et al. 1989; Sannier et al. 2002; Kogan et al. 2004; Wessels et al. 2006), MODIS (Kawamura et al. 2005; Xu et al. 2008), Landsat-5 TM and Landsat-7 ETM+ (Friedl et al. 1994; Schino et al. 2003; Samimi and Kraus 2004), SPOT VEGETATION (Verbesselt et al. 2006), and hyperspectral sensors such as PROBE-1, Hyperion, and the HyMap system (Mutanga and Skidmore 2004; Mirik et al. 2005; Numata et al. 2008; Cho and Skidmore 2009). Although many studies have investigated the ability to assess AGB from VI's, many problems have been found. One problem is that an empirical relationship derived by a VI for the accurate prediction of AGB at one site or time period may not apply to other sites or even the same site at another time (Foody et al. 2003). This problem is primarily due to variations in the natural environment (e.g., variable precipitation, soil-water content, and temperature conditions), viewing season (e.g., phenology during the growing season), and the sensor used in the study (e.g., differences in spatial resolution and other sensor characteristics) (Davidson and Csillag 2001; Schino et al. 2003; Flynn et al. 2008). In addition, since VI's have differing abilities to provide accurate estimates of AGB, it is difficult to determine an optimal VI for a specific study. For example, the same VI (e.g., NDVI) may have different prediction accuracies within various regions; yet different types of VI's (e.g., NDVI vs NDWI) may perform quite differently within the same region (Table 1). These problems limit the transferability of predictive relationships and the effectiveness of VI's to estimate AGB.

Approximately 48 percent of Idaho is considered rangeland, and many of these areas are categorized as a semiarid sagebrush-steppe ecosystem (<http://www.idrange.org>). AGB estimation in the semiarid rangelands of the Intermountain West plays an important role in rangeland ecosystem assessment. In the semiarid rangelands of Idaho, high temperatures hasten the desiccation of plants and many grass species senesce during the summer. The relationship between herbaceous AGB and VI's is most accurately estimated when the proportion of green or growing material is high (Hill 2004; Numata et al. 2008) relative to the proportion of bare ground and/or litter. While important, determining an optimal VI for the accurate estimation of seasonal herbaceous AGB in semiarid rangelands may be difficult.

Most VI's used for AGB estimation are based on radiance or reflectance from a red band (RED) around 0.66 μm and a near infrared band (NIR) around 0.86 μm (Huete et al. 2002; Chuvieco et al. 2004). The RED band characteristically shows a strong chlorophyll absorption region for vegetation and strong reflectance for soils, while the NIR band is located in the high reflectance plateau of vegetation canopies. Since absorption by liquid water near 0.86 μm is negligible, NIR reflectance is affected primarily by internal leaf structure and cellulose content (Gao 1996). In contrast, the short-wave infrared band (SWIR) (around 1.24 μm) is located in the high reflectance plateau of vegetation reflectance with weak liquid absorption (canopy scattering enhances the water absorption) (Jacquemoud et al. 1996; Jackson et al. 2004). The SWIR band reflects changes in both the vegetation water content and the spongy mesophyll structure of vegetation. The combination of the NIR band with the SWIR band can remove variation induced by internal leaf structure and leaf dry matter content (Gao 1996; Ceccato et al. 2001). This combination of these bands (NIR and SWIR) is also sensitive to changes in liquid water content within the vegetation canopy (Serrano et al. 2000; Zarco-Tejada et al. 2003).

In this study, a suite of water sensitive vegetation indices (WSVI) were developed incorporating the NIR and SWIR portions of the electromagnetic spectrum, to help characterize plant water content and better estimate herbaceous AGB in semiarid rangeland ecosystems. The study was designed to investigate the applicability of various VI's for the assessment of herbaceous AGB in the semiarid rangelands of Idaho, USA. To accomplish this, eight VI's including difference vegetation index (DVI, Richardson and Everitt 1992), ratio vegetation index (RVI, Jordan 1969), normalized difference vegetation index (NDVI, Rouse et al. 1973), re-normalized difference vegetation index (RDVI, Roujean and Breon 1995), soil adjusted vegetation index (SAVI, Huete 1988), the second modified soil adjusted vegetation index (MSAVI2, Qi et al. 1994), infrared percentage vegetation index (IPVI, Crippen 1990), and modified simple ratio (MSR, Chen 1996), were derived from Satellite Pour l'Observation de la Terre 5 (SPOT 5) imagery. In addition, the relationship between herbaceous AGB and total water content was determined. Finally, correlation estimates between herbaceous AGB, VI's, and WSVI's were calculated, and the performance of herbaceous AGB predictions from both VI's and WSVI's were evaluated using field-based measurements of herbaceous AGB.

MATERIALS AND METHODS

Study Area

The study area, known as the Big Desert, lies in southeast Idaho, USA, approximately 71 km northwest of Pocatello. The center of the study area was located at 113° 4' 18.68" W and 43° 14' 27.88" N (Figure 1). This area is managed by the US Bureau of Land Management (BLM) and exhibits a large variety of native as well as invasive plant species. The area is a semiarid sagebrush-steppe ecosystem with a high proportion of bare ground (\bar{x} bare ground > 17%, Studley et al. 2009). The area is sagebrush-steppe, consisting primarily of native and non-native grasses, forbs, and many shrub species including sagebrush (*Artemisia tridentata*) and rabbit brush (*Chrysothamnus nauseosus*). Annual precipitation is 23 cm with 40% of the precipitation falling from April through June. The area is bordered by geologically young lava formations to the south and west and irrigated agricultural lands to the north and east. Sheep grazing is the primary anthropogenic disturbance to the study area with semi-extensive continuous/seasonal grazing systems used on allotments ranging in size from 1100 to over 125,000 ha. Wildfire is a common disturbance and nearly 40% of the study area has burned in the past 10 years.




Figure 1. Location and general characteristics of the Big Desert in southeastern, Idaho. Note: no weather station survey site was available within the Big Desert study area however, nine sites were located which bound the study area. Though some sites are in the mountains, the weather there has identical change trends compared to the snake river plain.

Field Data Collection

This study presents results using total herbaceous AGB measurements only and does not include any measurements of shrub biomass production. Twenty-nine sample locations were selected for the collection of herbaceous AGB, which has been defined as all grasses, forbs, and standing litter for the purposes of this study. Site selection criteria included the site being a homogeneous area at least 20 m x 20 m in size (cf., spatial resolution of SPOT satellite imagery = 10 m x 10 m in size, thus helping to assure the sample pixel was also homogeneous) with still larger areas being preferred. The domination plants in each site are herbaceous vegetations, with the plot center ≥ 70 meters from any “edges” including roads, fences, or power lines, and plot perimeters ≥ 100 meters from all other plots. Preference was given to sites with perimeters located >250 meters apart. The location of each sample plot center was recorded using a Trimble Geo XH GPS receiver using latitude-longitude (WGS 84). All GPS data were post-process differentially corrected (± 0.10 m after post processing with a 95% CI using reference stations located <80 km from the study area) to ensure the sample location was registered with the correct and representative pixel within the satellite imagery (Weber 2006; Weber et al. 2008).

Available herbaceous AGB was measured using a plastic coated cable hoop 2.36 meters in circumference. The hoop was randomly tossed into each of four quadrants (NW, NE, SE, and SW) centered over the sample point. All herbaceous vegetation within the hoop were clipped as close to the ground as allowed by the clipper (approximately 5mm of the ground surface) and weighed immediately (± 1 g) using a Pesola scale tared to the weight of an ordinary paper bag. The samples were taken to the laboratory, and dried in 75 °C ovens for 48 hours. After drying, the samples were re-weighed to determine vegetation water content. Biomass was estimated following Sheley (1999) and expressed in kilograms per hectare.

Vegetation Indices Derived from SPOT 5 Imagery

Satellite Pour l'Observation de la Terre 5 (SPOT 5) multispectral imagery (10 m x 10 m pixels) was acquired for the Big Desert study area on June 27, 2009. The imagery was georectified against 2004 National Agriculture Imagery Program (NAIP) natural color aerial imagery (1 m x 1 m pixels).

Atmospheric correction was performed with Idrisi Taiga (v16.03) using the ATMOSC module (Clark Labs, Worcester, MA). All imagery was corrected for atmospheric effects using the Cos(t) model (Chavez 1996) and input parameters reported in the metadata supplied by SPOT Image Corporation. The imagery was then projected into Idaho Transverse Mercator (NAD 83). The eight VI's used in this study were derived from the SPOT 5 imagery.

Water Sensitive Vegetation Indices

Using the same SPOT5 imagery, eight WSVI's were developed by directly substituting the SWIR band for the RED band within the eight VI's described above (Table 2). The "Sample" tool within ESRI's ArcGIS 10 was then used to extract VI and WSVI values at each sample site ($n = 29$). The resulting data were exported to SPSS (V17.0) for further analysis. Correlations between VI/WSVI values and measured herbaceous AGB were used to determine the applicability and efficacy of each.

Table 2. Water sensitive vegetation indices (WSVI) used to estimate herbaceous total above ground biomass (AGB) (note the substitution of the SWIR band for the RED band [cf. table 2])

<i>Index</i>	<i>Formula</i>
DWI	$NIR - SWIR$
RWI	$NIR/SWIR$
NDWI	$\frac{NIR - SWIR}{NIR + SWIR}$
RDWI	$\frac{NIR - SWIR}{\sqrt{NIR + SWIR}}$
SAWI	$\frac{(NIR - SWIR)(1 + L)}{NIR + SWIR + L}$, where $L = 0.5$
MSAWI2	$\frac{2NIR + 1 - \sqrt{(2NIR + 1)^2 - 8(NIR - SWIR)}}{2}$
IPWI	$\frac{NIR}{NIR + SWIR}$
MWSR	$\frac{\frac{NIR}{SWIR} - 1}{\sqrt{\frac{NIR}{SWIR} + 1}}$

RESULTS AND DISCUSSION

Field-based herbaceous AGB estimates ranged from 518 kg/ha to 8075 kg/ha ($\bar{x} = 2982$ kg/ha) based on vegetation samples collected at 29 field locations. Using linear regression analysis between each VI and herbaceous AGB measurements, the relationship between these variables was described (Table 3). Based upon these results, it was noted that the relationships varied greatly and the strength of all correlations were relatively weak ($0.28 \leq r^2 \leq 0.40$). This was likely attributable to the mixture of photosynthetic and non-photosynthetic plant material found in the field and correspondingly, in the herbaceous AGB samples used in this study. As a result, the VI's provided poor estimates of herbaceous AGB. Furthermore, the prediction of herbaceous AGB was least well explained using NDVI ($r^2 = 0.28$, $p = 0.003$) and as a result, NDVI was not considered a reliable predictor of herbaceous AGB in this study area, though it remains one of most widely used VI's for AGB prediction and many other vegetation studies.

Table 3. Correlation between herbaceous total above ground biomass (AGB) and the VI's and WSVI's used in this study (r^2 = coefficient of determination)

Standard VI's using Red and NIR bands				Water sensitive VI's using NIR and SWIR bands			
Index	r^2	F-value	p	Index	r^2	F-value	p
DVI	0.40	17.9	<0.001	DWI	0.53	30.1	<0.001
RVI	0.35	14.3	0.001	RWI	0.54	31.2	<0.001
NDVI	0.28	10.6	0.003	NDWI	0.52	29.0	<0.001
RDVI	0.35	14.3	0.001	RDWI	0.52	29.8	<0.001
SAVI	0.37	15.8	<0.001	SAWI	0.53	29.8	<0.001
MSAVI2	0.39	17.0	<0.001	MSAWI2	0.53	30.3	<0.001
IPVI	0.28	10.6	0.003	IPWI	0.52	29.0	<0.001
MSR	0.32	12.6	0.001	MWSR	0.53	30.3	<0.001

Based on field survey data, the relationship between herbaceous AGB and vegetation water content (Figure 2) revealed a significant correlation ($r^2 = 0.94$, $P < 0.001$). Related studies have shown that grass biophysical parameters such as leaf area index are related to liquid water content (Hunt and Rock 1989; Roberts et al. 1997, 2004). Numata et al. (2008) indicated water absorption spectra between 1100 and 1250 nm had a significant correlation with canopy water content and suggested the use of water absorption features (i.e., water absorption depth and water absorption area) may improve the accuracy of biomass estimation. Therefore, the hypothesis that an index that closely correlates to water content may also exhibit strong correlation with herbaceous AGB was tested.




Figure 2. Relationship between herbaceous total above ground biomass (AGB) and vegetation water content.

In order to validate this hypothesis, the eight VI's used in this study along with eight WSVI's were correlated against vegetation water content (Table 4). The observed correlation between the VI's and vegetation water content were relatively weak ($0.30 \leq r^2 \leq 0.40$), while the WSVI's were more highly correlated ($r^2 \geq 0.51$) with vegetation water content. These results suggest the combination of NIR and SWIR bands are more sensitive to changes in liquid water content within vegetation, and that WSVI's exhibit a better response to water content of herbaceous vegetation in semiarid rangeland ecosystems.

Table 4. Coefficients of determination (r^2) calculated for the relationships between vegetation water content and vegetation indices

Standard VI's using Red and NIR bands				Water sensitive VI's using NIR and SWIR bands			
<i>Index</i>	r^2	<i>F-value</i>	<i>p</i>	<i>Index</i>	r^2	<i>F-value</i>	<i>p</i>
DVI	0.40	17.9	<0.001	DWI	0.52	29.4	<0.001
RVI	0.35	14.8	0.001	RWI	0.52	29.7	<0.001
NDVI	0.30	11.6	0.002	NDWI	0.52	28.5	<0.001
RDVI	0.36	15.0	0.001	RDWI	0.52	29.1	<0.001
SAVI	0.38	16.2	<0.001	SAWI	0.52	29.2	<0.001
MSAVI2	0.39	17.2	<0.001	MSAWI2	0.52	29.5	<0.001
IPVI	0.30	11.6	0.002	IPWI	0.51	28.5	<0.001
MSR	0.33	13.4	0.001	MWSR	0.52	29.2	<0.001

Linear relationships were determined between eight WSVI's and herbaceous AGB (Table 3). In comparison to the more traditional VI's ($r^2 \leq 0.40$), the WSVI's exhibited much stronger relationships with herbaceous AGB ($r^2 \geq 0.52$). In addition, it is noted that the NDWI (based on the same simple band ratio structure as NDVI [NDVI=(NIR-R)/(NIR+R)]), better explained the variation in herbaceous AGB relative to NDVI (i.e., coefficient of determination increased from 0.28 to 0.52). This result is similar to previous research reporting that NDWI performed better in drought conditions than NDVI (Verbesselt et al. 2006a; Gu et al. 2007). These results further suggest that herbaceous AGB is highly correlated with vegetation water content and that WSVI's can more accurately predict herbaceous AGB for semiarid rangeland sites.

This study was designed for herbaceous AGB estimation in semiarid rangeland of Idaho. The relationships revealed by the study are still condition-specific and should not be directly extrapolated to other regions, however, the specific approach developed in this study can be used across other rangeland areas.

Assessment of Error and Bias

It is difficult to collect a large numbers of field-measured AGB data, and we used a relatively small sample size ($n=29$) in this study. Because the limitation on sample size may influence the strength of each index, the robustness of the correlation between WSVI's and AGB were tested using the jackknife method (Table 5) (Efron and Gong 1983; Buermann et al. 2008). Small RMSE values were computed and the mean r-square values calculated by the jackknife method were similar to the r-square values given in Table 3. We conclude that the observed correlation between WSVI's and AGB were not highly influenced by a few individual samples and the regression and correlation results presented herein are robust.

Table 5. The robustness of the correlation between WSVI's and AGB were tested by the jackknife method

<i>Index</i>	<i>Mean r^2</i>	<i>RMSE</i>
DWI	0.52	0.032
RWI	0.54	0.026
NDWI	0.51	0.038
RDWI	0.52	0.031
SAWI	0.51	0.036
MSAWI2	0.51	0.038
IPWI	0.52	0.019
MWSR	0.53	0.037

Previous studies have demonstrated varied results with VI's and each reveal different strengths of correlations with AGB under specific conditions dependent upon the phenology of plants within a given growing season (Reeves et al. 2001). Cho and Skidmore (2009) indicate VI's are highly correlated ($r^2 \geq 0.50$) with AGB when the vegetation was in the early stages of senescence. In semiarid rangeland ecosystems, high summer temperatures hasten the desiccation of plants, and many plants begin senescence in mid-late June. In this study, all herbaceous AGB data were collected between July 1 and July 9, 2009. Based on monthly precipitation data provided by the United States Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS) (<http://www.id.nrcs.usda.gov/snow/data/historic.html>) and the United States Bureau of Reclamation (USBR) AgriMet Program (<http://www.usbr.gov/pn/agrimet/>), it is noted that mean monthly precipitation between April and July 2009 (260 mm) was greater than the monthly precipitation during the same time period in either 2007 (140 mm) or 2008 (92 mm), and that precipitation in June substantially increased in 2009 (48 mm [2007], 21mm [2008], 141 mm [2009]). (Figure 1) (Table 6; Table 7). These statistics, along with cooler than average temperatures, suggest that senescence may have been delayed in 2009, with the experimental time period of this study falling within the early stages of senescence. Numata et al. (2008) achieved very poor correlation between NDWI/NDVI and herbaceous AGB (Table 1) because their field sampling occurred much later in the season (beginning of August) and during a time of the year when most herbaceous plant materials were already senesced.

Table 6. Natural Resources Conservation Service (NRCS) and AgriMet survey site list along with monthly precipitation (mm) and monthly mean temperature (°C)

Site name	Lat	Long	Year	Precipitation					Mean temperature		
				April	May	June	July	Total	May	June	July
Garfield R.S.	43°36'	-113°55'	2007	43	8	25	28	104	9	13	19
			2008	23	43	13	23	102	16	19	20
			2009	30	71	203	20	324	8	10	16
Swede Peak	43°37'	-113°58'	2007	56	15	30	38	139	8	13	19
			2008	25	28	3	0	56	7	12	18
			2009	28	43	201	23	295	7	9	15
Smiley Mountain	43°43'	-113°50'	2007	89	33	89	15	226	6	10	17
			2008	30	94	48	13	185	3	8	15
			2009	89	91	180	43	403	5	7	13
Howell Canyon	42°19'	-113°36'	2007	150	30	117	25	322	8	12	18
			2008	56	76	58	15	205	5	10	17
			2009	124	66	170	3	363	7	9	16
Wildhorse Divide	42°45'	-112°28'	2007	86	25	74	13	198	10	13	18
			2008	36	66	33	0	135	7	12	17
			2009	107	58	163	36	364	9	11	16
Fort Hall	43°04'	-112°25'	2007	30	5	39	5	79	13	17	23
			2008	3	37	9	4	53	11	16	20
			2009	31	29	97	15	172	12	15	20
Rupert	42°35'	-113°52'	2007	19	9	20	8	56	14	17	23
			2008	3	17	12	7	39	12	17	21
			2009	26	18	53	6	103	13	15	21
Picabo	43°18'	-114°09'	2007	23	3	10	3	39	13	18	24
			2008	7	22	7	0	36	11	16	21
			2009	20	41	98	8	167	12	14	20
Aberdeen	42°57'	-112°49'	2007	36	3	30	14	83	14	18	23
			2008	4	23	6	1	34	12	16	21
			2009	23	22	102	17	164	13	15	20

Table 7. Analysis of precipitation (mm) and temperature (°C) on 2007, 2008 and 2009

Year	Average precipitation				Average temperature			Total precipitation	Standard deviation
	April	May	June	July	May	June	July		
2007	59	15	48	17	11	15	20	140	56 ¹
2008	20	45	21	7	9	14	19	92	
2009	53	49	141	19	10	12	17	260	69 ²

¹Standard deviation of precipitation for 2007 and 2008.

²Standard deviation of precipitation for 2008 and 2009.

NDVI physically responds to chlorophyll absorption and is not directly related to the quantity of water in the vegetation (Ceccato et al. 2002). Cheng et al. (2008) indicate that NDVI shows correlation with water content was probably due to the correlation to green leaf density. Because of the sparse vegetation and dry plant matter (litter) found in semiarid regions, NDVI was not a reliable indicator of water content or herbaceous AGB in this study. However, some AGB estimates have shown strong NDVI correlations to biomass ($r^2 \geq 0.50$), but this may be because these study areas were more homogeneous and/or contained a higher proportion of green grass cover (Mašková et al. 2008). The accuracy of herbaceous AGB predictions based upon remotely sensed data is strongly influenced by the presence and abundance of grass species as well as the presence and abundance of bare ground and other spectral distraction features. A more homogeneous surface always provides higher correlations between remotely sensed measures and herbaceous AGB estimates compared to more heterogeneous surfaces (Numata et al. 2008). In addition, as opposed to the traditional VI, the WSVI's have the advantage of leveraging liquid water absorption regions to more accurately predict water content even in areas without contiguous spectral coverage (Serrano et al. 2000). This is possibly one reason why the WSVI's performed better in the semiarid rangelands of Idaho.

CONCLUSION

This study focused on the estimation of herbaceous AGB in the semiarid rangelands of Idaho. Based on a survey of herbaceous AGB, a significant correlation ($p < 0.001$) between herbaceous AGB and vegetation water content was found. In addition, a suite of WSVI's were developed that describe water content and herbaceous AGB in semiarid rangeland ecosystems. Correlation estimates between herbaceous AGB, VI's, and WSVI's were calculated, and the performance of herbaceous AGB predictions for both the VI's and WSVI's were evaluated using field-based measurements of herbaceous AGB. Results demonstrate the WSVI's were correlated ($r^2 \geq 0.51$) with vegetation water content and performed better in herbaceous AGB estimation for the semiarid rangelands of Idaho relative to VI's. Furthermore, it was noticed that not only did vegetation water content influence the accuracy of herbaceous AGB estimates, but based on findings reported in other studies, phenological stage and plant community structure also influence the accuracy of herbaceous AGB estimates derived from remotely sensed data. Numerous factors influence the successful use of remote sensing data for the estimation of herbaceous AGB and water content, described using WSVI's, explained approximately 50% of the variance in herbaceous AGB measurements collected as part of this study. Other factors that likely play a role include sun angle, shadow, georegistration, and the varying affect of soils. Future work will seek to assess a more comprehensive characterization of the influence of these factors on herbaceous AGB estimations in semiarid rangelands.

ACKNOWLEDGMENTS

This study was made possible by a grant from the National Aeronautics and Space Administration Goddard Space Flight Center (NNX08AO90G). Idaho State University would also like to acknowledge the Idaho Delegation for their assistance in obtaining this grant.

LITERATURE CITED

Boelman, N.T., M. Stieglitz, H.M. Rueth, M. Sommerkorn, K.L. Griffin, G.R. Shaver, and J.A. Gamon, 2003. Response of NDVI, Biomass, and Ecosystem Gas Exchange to Long-term Warming and Fertilization in Wet Sedge Tundra, *Oecologia*, 135(3): 414-421

- Box, E.O., B.N. Holben, and V. Kalb, 1989. Accuracy of the AVHRR Vegetation Index as a Predictor of Biomass, Primary Productivity and Net CO₂ Flux, *Plant Ecology*, 80(2): 71-89
- Breman, H., and C.T. de Wit, 1983. Rangeland Productivity and Exploitation in the Sahel, *Science*, 221(4618): 1341-1347
- Buermann, W., S. Saatchi, T.B. Smith, B.R. Zutta, J.A. Chaves, B. Milá, and C.H. Graham, 2008. Predicting Species Distributions Across the Amazonian and Andean Regions Using Remote Sensing Data, *Journal of Biogeography*, 35(7): 1160-1176
- Ceccato, P., S. Flasse, S. Tarantola, S. Jacquemond, and J.M. Gregoire, 2001. Detecting Vegetation Leaf Water Content Using Reflectance in the Optical Domain, *Remote Sensing of Environment*, 77(1): 22-33
- Ceccato, P., S. Flasse, and J.M. Gregoire, 2002. Designing a Spectral Index to Estimate Vegetation Water Content from Remote Sensing Data: Part 2. Validation and Applications, *Remote Sensing of Environment*, 82(2-3): 198-207
- Chavez, P.S., 1996. Image-based Atmospheric Corrections: Revisited and Improved, *Photogrammetric Engineering and Remote Sensing*, 62(9): 1025-1036
- Chen, J.M., 1996. Evaluation of Vegetation Indices and a Modified Simple Ratio for Boreal Applications, *Canadian Journal of Remote Sensing*, 22(3): 229-242
- Cheng, Y.B., S.L. Ustin, D. Riaño, and V.C. Vanderbilt, 2008. Water Content Estimation from Hyperspectral Images and MODIS Indexes in Southeastern Arizona, *Remote Sensing of Environment*, 112(2): 363-374
- Cho, M.A., and A.K. Skidmore, 2009. Hyperspectral Predictors for Monitoring Biomass Production in Mediterranean Mountain Grasslands: Majella National Park, Italy, *International Journal of Remote Sensing*, 30(2): 499-515
- Chuvieco, E., D. Cocero, D. Riaño, P. Martin, J. Martínez-Vega, J.D.L. Riva, and F. Pérez, 2004. Combining NDVI and Surface Temperature for the Estimation of Live Fuels Moisture Content in Forest Fire Danger Rating, *Remote Sensing of Environment*, 92(3): 322-331
- Crippen, R.E., 1990. Calculating the Vegetation Index Faster, *Remote Sensing of Environment*, 34(1): 71-73
- Davidson, A., and F. Csillag, 2001. The Influence of Vegetation Index and Spatial Resolution on a Two-date Remote Sensing Derived Relation to C4 Species Coverage, *Remote Sensing of Environment*, 75(1): 138-151

- Efron, B., and G. Gong, 1983. A Leisurely Look at the Bootstrap, the Jackknife, and Cross-validation, *American Statistician*, 37(1): 36-48
- Flynn, E.S., C.T. Dougherty, and O. Wendroth, 2008. Assessment of Pasture Biomass with the Normalized Difference Vegetation Index from Active Ground-based Sensors, *Agronomy Journal*, 100(1): 114-121
- Foody, G.M., D.S. Boyd, and M.E.J. Cutler, 2003. Predictive Relations of Tropical Forest Biomass from Landsat TM Data and Their Transferability between Regions, *Remote Sensing of Environment*, 85(4): 463-474
- Friedl, M.A., J. Michaelsen, F.W. Davis, H. Walker, and D.S. Schimel, 1994. Estimating Grassland Biomass and Leaf Area Index Using Ground and Satellite Data, *International Journal of Remote Sensing*, 15(7): 1401-1420
- Gao, B.C, 1996. NDWI : A Normalized Difference Water Index for Remote Sensing of Vegetation Liquid Water from Space, *Remote sensing of environment*, 58(3): 257-266
- Gu, Y., J.E. Brown, J.P. Verdin, and B. Wardlow, 2007. A Five-year Analysis of MODIS NDVI and NDWI for Grassland Drought Assessment over the Central Great Plains of the United States, *Geophysical Research Letter*, 34(L06407), doi:10.1029/2006GL029127
- Hill, M.J., 2004. Grazing Agriculture: Managed Pasture, Grassland, and Rangeland, Pages 449-530 in S. L. Ustin (Ed.) Manual of Remote Sensing volume 4: Remote sensing for natural resource management and environmental monitoring. Hoboken, NJ, United States: John Wiley & Sons.
- Huete, A.R., 1988. A Soil-adjusted Vegetation Index (SAVI), *Remote sensing of environment*, 25(3): 295-309
- Huete, A.R., K. Didan, T. Miura, E.P. Rodriguez, X. Gao, and L.G. Ferreira, 2002. Overview of the Radiometric and Biophysical Performance of the MODIS Vegetation Indices, *Remote Sensing of Environment*, 83(1-2): 195-213
- Hunt, E.R. and B.N. Rock, 1989. Detection of Changes in Leaf Water Content Using Near- and Middle-Infrared Reflectances, *Remote Sensing of Environment*, 30(1): 43-54
- Huntsinger, L. and P. Hopkinson, 1996. Viewpoint: Sustaining Rangeland Landscapes: A Social and Ecological Process, *Journal of Range Management*, 49(2): 167-173
- Jackson, T.J., D. Chen, M. Cosh, F. Li, M. Anderson, C. Walthall, P. Doriaswamy, and E.R. Hunt, 2004. Vegetation Water Content Mapping Using Landsat Data Derived Normalized Difference Water Index for Corn and Soybeans, *Remote Sensing of Environment*, 92(4): 475-482

- Jacquemoud, S., S.L. Ustin, J. Verdebout, G. Schmuck, G. Andreoli, and B. Hosgood, 1996. Estimating Leaf Biochemistry Using the PROSPECT Leaf Optical Properties Model, *Remote Sensing of Environment*, 56(3): 194-202
- Jordan, C.F., 1969. Derivation of Leaf Area Index from Quality of Light on the Forest Floor, *Ecology*, 50(4): 663-666
- Kawamura, K., T. Akiyama, H. Yokota, M. Tsutsumi, T. Yasuda, O. Watanabe, and S. Wang, 2005. Comparing MODIS Vegetation Indices with AVHRR NDVI for Monitoring the Forage Quantity and Quality in Inner Mongolia Grassland, China, *Grassland Science*, 51(1): 33-40
- Kogan, F., R. Stark, A. Gitelson, L. Jargalsaikhan, and S. Tsooj, 2004. Derivation of Pasture Biomass in Mongolia from AVHRR-based Vegetation Health Indices, *International Journal of Remote Sensing*, 25(14): 2889-2896
- Masková, Z., F. Zemek, and J. Kvet, 2008. Normalized Difference Vegetation Index (NDVI) in the Management of Mountain Meadows, *Boreal Environmental Research*, 13(5): 417-432
- Mirik, M., J. Norland, R. Crabtree, and M. Biondini, 2005. Hyperspectral One-meter-resolution Remote Sensing in Yellowstone National Park, Wyoming: II. Biomass, *Rangeland Ecology and Management*, 58(5): 459-465
- Mutanga, O., and A.K. Skidmore, 2004. Hyperspectral Band Depth Analysis for a Better Estimation of Grass Biomass (*Cenchrus ciliaris*) Measured under Controlled Laboratory Conditions, *International Journal of Applied Earth Observation and Geoinformation*, 5(2): 87-96
- Numata, I., D.A. Roberts, O.A. Chadwick, J.P. Schimel, L.S. Galvão, and J.V. Soares, 2008. Evaluation of Hyperspectral Data for Pasture Estimate in the Brazilian Amazon Using Field and Imaging Spectrometers, *Remote Sensing of Environment*, 112(4): 1569-1583
- Qi, J., A. Chehbouni, A.R. Huete, Y.H. Kerr, and S. Sorooshian, 1994. A Modified Soil Adjusted Vegetation Index. *Remote Sensing of Environment*, 48(2): 119-126
- Reeves, M.C., J.C. Winslow, and S.W. Running, 2001. Mapping Weekly Rangeland Vegetation Productivity Using MODIS Algorithms, *Rangeland Ecology & Management*, 54: A90-A105
- Richardson, A.J., and J.H. Everitt, 1992. Using Spectral Vegetation Indices to Estimate Rangeland Productivity. *Geocarto International*, 7(1): 63-69
- Roberts, D.A., R.O. Green, and J.B. Adams, 1997. Temporal and Spatial Patterns in Vegetation and Atmospheric Properties from AVIRIS. *Remote Sensing of Environment*, 62(3): 223-240

- Roberts, D.A., S.L. Ustin, S. Ogunjemiyo, J. Greenberg, S.Z. Dobrowski, J.Q. Chen, and T.M. Hinckley, 2004. Spectral and Structural Measures of Northwest Forest Vegetation at Leaf to Landscape Scales, *Ecosystems*, 7(5): 545-562
- Roujean, J.L., and F.M. Breon, 1995. Estimating PAR Absorbed by Vegetation from Bidirectional Reflectance Measurements, *Remote Sensing of Environment*, 51(3): 375-384
- Rouse, J.W., R.H. Haas, J.A. Schell, and D.W. Deering, 1973. Monitoring Vegetation Systems in the Great Plains with ERTS, in the 3rd Earth Resources Technology Satellite Symposium, 10-14 December 1973, Washington, D.C., USA, 309-317
- Running, S.W., R.R. Nemani, F.A. Heinsch, M. Zhao, M.C. Reeves, and H. Hashimoto, 2004. A Continuous Satellite-derived Measure of Global Terrestrial Primary Production, *BioScience*, 54(6): 547-560
- Samimi, C., and T. Kraus, 2004. Biomass Estimation Using Landsat-TM and -ETM+. Towards a Regional Model for Southern Africa? *GeoJournal*, 59(3): 177-187
- Sannier, C.A.D., J.C. Taylor, and W. Du Plessis, 2002. Real-time Monitoring of Vegetation Biomass with NOAA-AVHRR in Etosha National Park, Namibia, for Fire Risk Assessment, *International Journal of Remote Sensing*, 23(1): 71-89
- Schino, G., F. Borfecchia, L. Cecco, C. Dibari, M. Iannetta, S. Martini, and F. Pedrotti, 2003. Satellite Estimate of Grass Biomass in a Mountainous Range in Central Italy, *Agroforestry Systems*, 59(2): 157-162
- Serrano, L., S.L. Ustin, D.A. Roberts, J.A. Gamon, and J. Peñuelas, 2000. Deriving Water Content of Chaparral Vegetation from AVIRIS Data, *Remote Sensing of Environment*, 74(3): 570-581
- Sheley, R. 1999. AUM Analyzer Software, Montana State University, Bozeman, Mont. URL = <http://www.montana.edu>, visited 31-March-2010
- Studley, H. and K.T. Weber, 2009. Range Vegetation Assessment in the Big Desert, Upper Snake River Plain, Idaho, URL = http://giscenter.isu.edu/research/Techpg/nasa_postfire/To_PDF/2009_Big_Desert_FieldReport.pdf visited 31-March-2010
- Tucker, C.J., 1979. Red and Photographic Infrared Linear Combinations for Monitoring Vegetation, *Remote Sensing of Environment*, 8(2): 127-150
- Verbesselt, J., P. Jonsson, S. Lhermitte, J. van Aardt, and P. Coppin, 2006a. Evaluating Satellite and Climate Data-derived Indices as Fire Risk Indicators in Savanna Ecosystems, *IEEE Transactions on Geoscience and Remote Sensing*, 44(6): 1622-1632

Verbesselt, J., B. Somers, J. van Aardt, I. Jonckheere, and P. Coppin, 2006. Monitoring Herbaceous Biomass and Water Content with SPOT VEGETATION Time-series to Improve Fire Risk Assessment in Savanna Ecosystems, *Remote Sensing of Environment*, 101(3): 399-414

Weber, K.T., 2006. Challenges of Integrating Geospatial Technologies into Rangeland Research and Management, *Rangeland Ecology and Management*, 59(1): 38-43

Weber, K.T., J. Theau, and K. Serr, 2008. Effect of Co-registration Error on Patchy Target Detection Using High-resolution Imagery, *Remote Sensing of the Environment*, 112(3): 845-850

Weiser, R.L., G. Asrar, G.P. Miller, and E.T. Kanemasu, 1986. Assessing Grassland Biophysical Characteristics from Spectral Measurements, *Remote Sensing of the Environment*, 20(2): 141-152

Wessels, K.J., S.D. Prince, N. Zambatis, S. MacFadyen, P.E. Frost, and D. Van Zyl, 2006. Relationship between Herbaceous Biomass and 1-km² Advanced Very High Resolution Radiometer (AVHRR) NDVI in Kruger National Park, South Africa, *International Journal of Remote Sensing*, 27(5): 951-973

Xu, B., X.C. Yang, W.G. Tao, Z.H. Qin, H.Q. Liu, J.M. Miao, and Y.Y. Bi, 2008. MODIS-based Remote Sensing Monitoring of Grass Production in China, *International Journal of Remote Sensing*, 29(17): 5313-5327

Zarco-Tejada, P.J., C.A. Rueda, and S.L. Ustin, 2003. Water Content Estimation in Vegetation with MODIS Reflectance Data and Model Inversion Methods, *Remote Sensing of Environment* 85(1): 109-124

Recommended citation style:

Chen, F., K.T. Weber, B. Gokhale, 2011. Herbaceous Biomass Estimation from SPOT-5 Imagery in Semiarid Rangelands of Idaho. Pages 89-104 in K. T. Weber and K. Davis (Eds.), Final Report: Assessing Post-Fire Recovery of Sagebrush-Steppe Rangelands in Southeastern Idaho (NNX08AO90G). 252 pp.